



ISSN: 2456-2912

VET 2016; 1(3): 12-14

© 2016 VET

www.veterinarypaper.com

Received: 03-09-2016

Accepted: 04-10-2016

**Ashok Kumar Devarasetti**

Assistant Professor & Univ.  
Head, Department of Veterinary  
Biochemistry, College of  
Veterinary Science,  
Rajendranagar, Hyderabad,  
Telangana, India

**E Sunil Anand Kumar**

Assistant Professor & Principal,  
Department of Veterinary  
Gynaecology & Obstetrics,  
College of Veterinary Science,  
Rajendranagar, Hyderabad,  
Telangana, India

**K Venkata Ramana**

Associate Professor & Principal,  
Department of Veterinary  
Gynaecology & Obstetrics,  
College of Veterinary Science,  
Animal Husbandry Polytechnic,  
Mahaboobnagar, India

**L Ramsingh**

Assistant Professor, Department  
of Veterinary Gynaecology &  
Obstetrics, College of Veterinary  
Science, Rajendranagar,  
Hyderabad, Telangana, India

**Correspondence**

**Ashok Kumar Devarasetti**

Assistant Professor & Univ.  
Head, Department of Veterinary  
Biochemistry, College of  
Veterinary Science,  
Rajendranagar, Hyderabad,  
Telangana, India

## Supplementation of dietary yeast on body performance in Japanese quails

**Ashok Kumar Devarasetti, E Sunil Anand Kumar, K Venkata Ramana and L Ramsingh**

### Abstract

The objective of this study was to investigate the effect of supplementation of Dietary baker's yeast as a probiotic on body performance of Japanese quails (*Coturnix coturnix japonica*). In this experiment a total of 96 day-old unsexed, growing Japanese quails were equally divided into three groups of 32 birds in each group. The experiment was terminated when birds were 6 weeks old. The standard basal diet was fed in first group (G1), 5% and 10% level of baker's yeast was supplemented in basal diet for second group (G2) and third group (G3) respectively. Feed and water was supplied *ad libitum* for all the birds. Body performance was determined by measuring feed intake, body weight gain and feed conversion ration. The results obtained in this study showed that, supplementation of dietary baker's yeast had positive effect on production performance in Japanese quails.

**Keywords:** Japanese quails, dietary yeast, body performance, probiotics

### Introduction

Japanese quails (*Coturnix coturnix japonica*) have become an important live stock because it has a small body size, easy to handled, large number of birds can be kept in a limited space, high egg production, many offspring can be available from certain number of parents. It is also used in embryological studies (Ayasan and Okan, 2001) [1]. Quail eggs are rich in protein and good source of iron, phosphorus, riboflavin, and selenium (Bing, 2011) [2].

Probiotics are live microorganisms which are mainly derived from certain bacteria, fungi and yeast cell. Yeast (*Saccharomyces cerevisiae*) is one of the most widely used probiotics. It has been fed to animals and poultry. Yeast is a 'single cell protein' the production is originated in Germany during World War – I, when the baker's yeast, was grown with molasses as the carbon, energy and nitrogen source for consumption as a protein supplement.

It is known that, Mannan oligosaccharide and Fructo-oligosaccharides are derived from cell wall of yeast and shown in suppressing enteric pathogens and modulating the immune system in poultry (santini *et al.*, 2001; Spring *et al.*, 2000; Iji *et al.*, 2001) [10, 12, 5]. These properties led researchers to use yeast culture as probiotic feed additives in J. quails diet.

There was a limited number of studies reported on effect of dietary yeast on performance and blood biochemical profile in Japanese quails. Therefore, the present study was conducted to investigate the effect of supplementation of dietary baker's yeast as a probiotic on body performance and blood biochemical profile of Japanese quails

### Materials and Methods

In this study, a total of 96 laying Japanese quails aged 25 weeks were randomly assigned to 12 cages of 8 birds each and cages were allocated to three groups with four replication per treatment. The birds were housed in standard cages (40x40x25 cm<sup>3</sup>) in a temperature controlled house at 73 °F. All the birds had free access to feed and water. The photoperiod was 16 hrs (Vatsalya and Kashmiri, 2011) [13]. Prior to the supplementation of yeast in the experimental diets, all the birds were fed *ad libitum* for 7 days for acclimatization. The birds of three groups were fed with experimental diets as follows, In the first group (G<sub>1</sub>) was fed with control/basal diet (Table 1), in the second group (G<sub>2</sub>) was fed with 5% level of baker's yeast in the basal diet and in the third group (G<sub>3</sub>) was fed with 10% level of baker's yeast in the basal

diet. The experimental period lasted for 6 weeks.

**Table 1:** Ingredients and chemical composition of basal diet:

Ingredients	%
Ground yellow corn	57.83
Soya bean meal	32.94
Fish meal	3.50
Corn Gluten	3.48
Di calcium phosphate	0.33
Limestone	1.16
DL-Methionine	0.09
Lysine	0.07
Iodized sodium chloride	0.30
Minerals and vitamins premix	0.30
Calculated composition	
Crude protein (%)	24.0
Metabolic energy (Kcal/kg)	2900.0
Calorie/protein ratio (C/P)	120.83
Calcium (%)	0.80
Phosphorus (%)	0.30

**Table 2:** Effect of dietary treatments on performance of growing Japanese quails:

Item	Age in weeks	Treatments		
		Control group	5% yeast	10% yeast
Body weight gain		41.63 <sup>a</sup> ± 0.06	44.96 <sup>b</sup> ± 2.07	47.83 <sup>c</sup> ± 1.93
	0 to 2	77.31 <sup>a</sup> ± 0.03	84.37 <sup>b</sup> ± 1.05	95.74 <sup>c</sup> ± 2.23
	2 to 4	86.10 <sup>a</sup> ± 1.07	90.73 <sup>b</sup> ± 2.29	112.38 <sup>c</sup> ± 0.14
	4 to 6	204.94 <sup>a</sup> ± 1.16	219.96 <sup>b</sup> ± 5.41	255.95 <sup>c</sup> ± 4.30
	0 to 6			
Feed intake		40.51 <sup>a</sup> ± 1.24	43.04 <sup>b</sup> ± 2.25	45.42 <sup>c</sup> ± 1.26
	0 to 2	76.29 <sup>a</sup> ± 1.10	82.81 <sup>b</sup> ± 0.18	92.36 <sup>c</sup> ± 1.53
	2 to 4	85.23 <sup>a</sup> ± 0.38	88.14 <sup>b</sup> ± 1.17	99.81 <sup>c</sup> ± 0.39
	4 to 6	202.03 <sup>a</sup> ± 2.72	213.99 <sup>b</sup> ± 3.60	237.59 <sup>c</sup> ± 2.93
	0 to 6			
Feed conversion ratio		0.973 ± 2.35	0.957 ± 1.08	0.949 ± 1.53
	0 to 2	0.986 ± 1.06	0.981 ± 1.83	0.964 ± 0.68
	2 to 4	0.989 ± 2.86	0.971 ± 0.93	0.887 ± 2.78
	4 to 6	1.985 ± 2.34	0.972 ± 1.28	0.928 ± 0.68
	0 to 6			

In the present study, the data illustrated in table: 2 indicated that, bakers dried yeast supplementation significantly reduced feed intake, but the body weight gain was significantly higher, meanwhile feed conversion ratio showed significant improvement due to yeast supplementation. This positive enhancement in feed conversion efficiency was confirmed by Zeweil (1997) [14]; Chumpawadee *et al.*, (2009) [3]. The positive response on body weight gain as a result of adding yeast may be due to mannan oligosaccharides (MOS) from yeast cell walls (Newman and Newman, 2001; O'Quinn *et al.*, 2001) [6, 7]. Some research studies suggest that MOS may improve growth performance in young pigs (Davis *et al.*, 1999; Pettigrew, 2000) [4, 8] reported that supplemented yeast increased weight gain and feed conversion ration of broilers. The positive effect on production performance in quails may be due to the number of anaerobic and cellulolytic bacteria were increased when the experimental diet was supplemented with

The body weight (g/bird), body weight gain (g/bird) and feed intake (g) of birds per replicate were recorded on individual basis at weekly intervals and average values were recorded. Feed conversion ratio (FCR) was also calculated weekly. FCR was calculated by dividing the feed intake by body weight gain (Quigley *et al.*, 1997) [9].

Data were statistically analyzed using the analysis of variance. Significant difference between treatment means were calculated according to Snedecor and Cochran (1980) [11].

## Results and Discussion

The feed ingredients used in the control group and treatment group were same but 5% and 10% level of yeast was supplemented in group-2 and group-3 respectively. The effect of yeast culture on body weight gain, feed intake and feed conversion are shown in table: 2 It was observed that, yeast at levels of 5% and 10% to the basal diet improved ( $P>0.05$ ) body weight gain.

yeast which enhanced lactate utilization and moderate pH of gut, therefore, yeast improves the nutrients digestibility and growth performance revealed that dietary supplement of yeast (*S. cerevisiae*) improves the body performance in broilers. These above results may explain the significant effects of dietary yeast in improving the metabolic process.

## Conclusion

Dietary yeast have great potential to beneficially affect the gut microflora and hence improves the digestibility and health in Japanese quails. The present study confirms that, the supplementation of bakers yeast as a probiotic in growing Japanese quails diets significantly improved the body performance. It could be concluded that, dietary yeast to growing quails up to 10% level improved the performance and biochemical profile.

## References

1. Ayasan T, Okan F. The effect of diet with different probiotic (Protexin) levels on the fattening performance and carcass characteristics of Japanese quails. Proceedings of XV European symposium on the quality of poultry meat, Kubadasi, Turkey, 2001, 169-174.
2. Bing. free calorie and nutrition data information for egg, quail, whole, fresh, raw. View nutrition labels and sign up for a free online diet program, 2011. <http://caloriecount.about.com/calories-egg-quail-whole-fresh-i1140>.
3. Chumpawadee S, Chantiratikul A, Santaweek S. Effect of dietary inclusion of cassava yeast as probiotic source on egg production and egg quality of laying hens. Int. J. Poult. Sci. 2009; 8:195-199.
4. Devis ME, Maxwell CV, Kegley *et al.* Efficacy of mannan oligosaccharide (Bio Mos) addition at two levels of supplemental copper on performance and immunocompetence of early-weaned pigs. J. Anim. Sci. 1999; (Suppl. 1):63.
5. Iji PA, Saki AA, Tivey DR. International structure and function of broiler chickens on diets supplemented with a mannan oligosaccharide. J. Sci. Food Agric. 2001; 81:1186-1192.
6. Newman KE, Newman MC. Evaluation of mannan oligosaccharides on the microflora and immunoglobulin status of sows and piglet performance. J. Anim. Sci. 2001; (Suppl.1):189.

7. O'Quinn PR, Funderburke DW, Tibbetts GW. Effects of dietary supplementation with mannan oligosaccharides on sow and litter performance in a commercial production system. *J. Anim. Sci.* 2001; 79(Suppl. 1):212.
8. Pettigrew JE. Bio-Mos effects on pig performance: A review in: T.P. Lyons and K.S. Jacques (ed.) *Biotechnology in the feed industry: Proc. Alltech's 16<sup>th</sup> Symp.* University Press Loughborough, UK, 2000.
9. Quigley JD, Drewry JJ, Murray LM, Ivey SJ. Body weight gain, feed efficiency, and fecal scores of dairy calves in response to Galactosyl lactose or Antibiotics in milk replacers, *J. Dairy Sci.* 1997; 80:1751-1754.
10. Santin E, Maiorka A, Macai M, Grecco M, Sanchez JC, Okada TM. Performance and intestinal mucosa development of broiler chickens fed diets containing *Saccharomyces cerevisiae* cell wall. *J. Appl. Poult. Res.* 2001; 10:236-244.
11. Snedecor GW, Cochran WG. *Statistical Methods.* 7<sup>th</sup> ed. The Iowa State University Press. Ames, Iowa, 1980.
12. Spring P, Wenk C, Dawson KA, Newmann KE. The effects of dietary mannonoligosaccharides on cecal parameter and the concentrations of enteric bacteria in the ceca of salmonella challenged broiler chicks. *Poult. Sci.* 2000; 79:205-211.
13. Vatsalya V, Kashmiri LA. Association between body weight growth and selected physiological parameters in male Japanese quail. *Int. J. Poult. Sci.* 2011; 10:680-684.
14. Zeweil HS. Evaluation of using some feed additives in growing Japanese quail diets. *J. Agric. Sci. Mansoura Univ.* 1997; (11):3611-3622.