



ISSN: 2456-2912

VET 2022; 7(6): 49-52

© 2022 VET

[www.veterinarypaper.com](http://www.veterinarypaper.com)

Received: 10-10-2022

Accepted: 15-11-2022

**Adhona Bhajana Wijaya Negara**  
Animal Husbandry Study  
Program, Faculty of Science and  
Technology, Panca Byudi  
Development University Medan,  
Indonesia

**Andhika Putra**  
Animal Husbandry Study  
Program, Faculty of Science and  
Technology, Panca Byudi  
Development University Medan,  
Indonesia

**Corresponding Author:**  
**Adhona Bhajana Wijaya Negara**  
Animal Husbandry Study  
Program, Faculty of Science and  
Technology, Panca Byudi  
Development University Medan,  
Indonesia

## Potential carrying capacity of corn straw land for beef cattle in Tanjung Selamat village, Deli Serdang Regency

**Adhona Bhajana Wijaya Negara and Andhika Putra**

**DOI:** <https://doi.org/10.22271/veterinary.2022.v7.i6a.456>

### Abstract

The purpose of this study was to analyze the potential of the area in the form of visualization of the potential of corn straw as a local feed source, find out which areas could be used as beef cattle development centers and formulate a strategy for utilizing corn straw as a source of beef cattle feed. This research was conducted in Tanjung Selamat Village, Sunggal District, Deli Serdang Regency. The research took place for 3 months starting from June to August 2019. This research used a survey method (purposive sampling) / sampling based on certain objectives and considerations with the conditions for sampling, namely agricultural plants that are suitable for harvesting and have large harvest areas. The research results obtained from the production of agricultural waste, namely corn straw from the samples in Tanjung Selamat Village, Sunggal District as much as 2.20 kg, total DM 425.46 tons, carrying capacity 186.5 ST, capacity to increase the cattle population was obtained at 186.5 ST. Corn straw agricultural waste has not been given optimally to cattle, corn straw given without fermentation based on dry matter needs can accommodate 186.42 ST/harvest/year, based on crude protein needs can accommodate 135.99 ST/harvest/year, straw utilization optimally corn is still able to increase the livestock population of 186.5 ST.

**Keywords:** Analysis of carrying capacity, corn straw, Tanjung Selamat village

### 1. Introduction

Tanjung Selamat Village is part of the Medan Tuntungan District, Medan City which is directly adjacent to Deliserdang Regency, with an area of  $\pm$  330.5 Ha with the following boundaries: To the north it is bordered by Asam Kumbang Village, Medan Selayang District, to the South it is bordered by Kelurahan Namo Gajah/Kemenangan Tani, on the east it is bordered by Tanjung Sari Village, Medan Selayang District, on the west it is bordered by the Belawan River, Deli Serdang Regency, this village is a village which was originally an agrarian village where the people made their main livelihood as farmers. Based on the aspect of road infrastructure, especially roads across factories in transporting raw goods and finished goods, they have undergone improvements with asphalt to the entire village. These conditions show that in Tanjung Selamat Village there have been many changes, from the old condition which was originally a village with large agricultural land and the original inhabitants working as farmers are now dry land with various industrial buildings standing, thus making significant changes to livelihoods. Villagers. Changes in livelihoods caused by land conversion are important to study because changes in work orientation for the people of Tanjung Selamat Village can not only have positive impacts, but also negative impacts. The issue of land conversion is not only a threat to farmers, the physical environment, the economy, and the social environment in that place, but land conversion can have a broad effect, Corn crop forage waste can be given directly to livestock and can also be given in processed form, such as hay and silage. Processed forage waste can be stored for a long time to be used/consumed during the dry/dry season when field grass is difficult to obtain. The potential of this animal feed waste can be utilized to support the Deli Serdang Regency Government's program in the field of beef cattle development. In addition, the Provincial Government of North Sumatra also has a special program for the development of beef cattle. The trend of increasing meat consumption in Indonesia shows a significant increase, in line with population growth, increased income and economic status of the Indonesian people.

Several factors hindered the provision of forage, namely the change in the function of land which was previously used as a source of forage into residential land, land for food crops and industrial crops. In addition, in general the availability of forage is also influenced by climate, so that during the dry season there is a shortage of forage for livestock and vice versa during the rainy season the amount is abundant. To overcome the shortage of grass, one of which is the use of agricultural waste as feed [1]. Some of these problems are also faced by farmers/breeders in Tanjung Selamat Village, Sunggal District, Deli Serdang Regency.

Beef cattle as ruminants can consume forage in a day as much as 10-15 percent of their body weight. A cow weighing 350 kg requires consumption of 52.5 kg of grass (forage) per day. This large amount of forage needs must always be available continuously every day. Provision of grass (forage) to cattle can be given in fresh form and can also be in the form of processed feed in the form of hay and silage. Hay-shaped feed is feed derived from dried forage, usually made when forage production is abundant and can be stored for a long time for use during the dry/dry season when grass (forage) is difficult to obtain. Meanwhile, silage is processed feed derived from forage through an anaerobic preservation process (fermentation) which at the same time can also increase the digestibility of the feed itself. This study aims to analyze the potential carrying capacity of corn straw as animal feed for beef cattle farming with the research target to determine the carrying capacity of corn straw as feed for beef cattle in Tanjung Selamat Village, Sunggal District, Deli Tengah Regency.

## 2. Research Methods

### 2.1 Place and time of research

This research was conducted in Tanjung Selamat Village, Sunggal District, Deli Serdang Regency. The research lasted for 3 months starting from June to August 2022.

### 2.2 Tools and materials

The tools used in this study were sickle knives used to cut forage, plastic rope, scissors, wooden stakes, scroll meters, sacks, analytical scales, stationery, calculators and cameras. The material used in this research is corn waste. The material was taken from agricultural land in Tanjung Selamat Village, Sunggal District, Deli Serdang Regency.

### 2.3 Research methods

This study uses a survey method. The survey method used in this research is purposive sampling. Purposive sampling is a sampling method based on certain objectives and considerations from the researcher. With the conditions for sampling, namely agricultural crops that are suitable for harvesting and have a large harvest area. Sampling was carried out in accordance with the required sample requirements and the required sample size was not questioned. This method is used to obtain quantitative information to analyze problems regarding corn straw waste.

### 2.4 Data collection

The data used in this study consisted of primary data, namely data obtained from field surveys and secondary data obtained from related agencies or offices such as the Livestock Service Office, the Food Crops and Horticulture Service, the Regional Development Planning Agency and the Central Bureau of Statistics. Primary data is obtained from sampling from calculations by:

1. Determine the corn cropland as a sampling site by using the purposive sampling method and measuring plots around 2x1 m per stretch for careful analysis.
2. Prepare sampling equipment such as knives, sickles, sacks, plastic bags, scales, plastic straps, tape measure, stationery, calculators and cameras.
3. Sampling was carried out by cutting 10 cm of corn straw above the soil surface and then weighing it to determine the fresh weight of corn straw.
4. Each research plot took 1 stalk of corn straw and then chopped it for analysis in the laboratory.

## 2.5 Observed Parameters

### 2.5.1 Production of corn straw fresh weight

The production of fresh weight of corn straw is the total weight of corn straw after the production of corn is harvested and before it withers due to loss of water.

### 2.5.2 Corn straw dry weight production

Production of corn straw dry weight is the total weight of fresh corn straw after heating at 33.8 °C for 7 hours.

### 2.5.3 Corn straw crude protein

Corn straw crude protein is the amount of crude protein contained in feed ingredients obtained from the multiplication of nitrogen content with a conversion factor of 6.225 using the Kjeldahl method.

### 2.5.4 Corn straw feed carrying capacity

The carrying capacity of corn straw feed is calculated by knowing the production of corn straw BK divided by the number of livestock BK needs. Livestock requirement assuming that one ruminant livestock unit (1 ST) requires an average dry matter of 6.25 kg/day or 2281.25 kg/year and a crude protein requirement of 0.06 kg/day or 21.9 kg/year [2]

## 2.6 Data analysis

Data analysis of corn straw holding capacity including production of fresh weight, dry weight production, crude protein and feed carrying capacity of corn straw were analyzed descriptively.

## 3. Research result

### 3.1 Production of fresh corn straw

Corn straw production fresh based on the tiled snippets carried out are listed in table 1.

**Table 1:** Yields of Fresh Corn Straw Production in Tanjung Selamat Village, Deli Serdang District.

No	Fresh Straw Weight (Kg)	Straw Weight Dry (Kg)	Total Production (Kg)
1	2040	4100	6140
2	2700	3200	5900
3	3200	5900	9100
4	4100	6140	10240
5	3040	4820	7860
6	4420	5820	10240
7	2940	3800	6740
8	4180	6220	10400
9	3600	6040	9640
10	3140	4280	7420
Total	33360	50320	83680
Average	3336	5032	8368

Based on the data in table 3 above and the sample area data, the production of fresh corn is 16,680 kg/ha

### 3.2 Cattle capacity

The results of the recapitulation of the analysis of the capacity of cattle using corn straw obtained during the study are presented in table 5 as in Appendix 2.

**Table 2:** Recapitulation of Cattle Carrying Capacity Analysis in Tanjung Selamat Village, Deli Serdang District.

Analysis Parameters	Amount
<b>Corn Straw Production</b>	
Dry Material	425.46 tonnes
Crude protein	32.76 tonnes
<b>Analysis of Corn Straw Carrying Capacity</b>	
Dry Material	186.42 ST/ year
Crude protein	135.99 ST/ year
KPPTS analysis	26.5 st

### 3.3 Corn straw production

The results of calculating corn straw production obtained during the study are presented in table 3.

**Table 3:** Calculation Results of Corn Straw Production

Parameter	Total Production (Tons/ ha)
Dry Material	425.46
Crude protein	32.76

Explained that the dry matter obtained was 425.46 and the crude protein obtained was 32.76. Corn straw is the second largest production of agricultural waste which can be used as cattle feed after rice straw. Syamsu, *et al.* [3] added that the largest production of agricultural waste was rice straw 85.81%, corn straw 5.84%, peanut straw 2.84%, soybean straw 2.54%, cassava shoots 2.29% and cassava straw. Narrow 0.68%. The content of agricultural waste (corn straw) that has not been processed from the laboratory results of the Sei Putih Goat Research Workshop is 63.75% DM and 7.70% PK. The results of the nutritional content of corn straw given to livestock are influenced by the harvest period. Corn plants are harvested young, the water content of corn plants is high, conversely, the older the harvested corn, the less water it will contain [4]. Forages derived from corn plants are expected to be used as a substitute for grass for ruminant livestock. Corn forages in the form of stems and leaves (corn straw) which are still dry or fresh are chopped to be made into silage thereby increasing straw nutrition and livestock palatability [5].

Forage for livestock is not only natural grass or cultivated grass, but forage derived from agricultural waste. The research results obtained from the production of agricultural waste, namely corn straw from the samples in Tanjung Selamat Village, Sunggal District, amounted to 2.20 kg. The total production of BK of corn straw was 425.46 tons, this amount was lower when compared to the results of research [6] which was 8490.18 tons of BK. The low production of corn straw in Tanjung Selamat Village, which is located in the lowlands, is caused by the relatively higher ambient temperature (25.4°). High temperatures will cause avapotranspiration so that the percentage of water content in corn straw decreases because the plants tend to be dehydrated and result in a lower weight of corn straw in the lowlands [7]. added that high ambient temperatures will affect the water content of the leaves and will experience a water deficit. This is consistent with observations when sampling corn straw at

the research location, corn straw seems to experience aging/drying faster so that the weight of corn straw is lower.

Rainfall factor also greatly affects the low production of corn straw [8]. Stated that normal rainfall for the growth of corn plants is 200-250 mm per month. Dry matter production depends on the availability of nutrients in the soil. Even though the same fertilizer was given at the study site, the effectiveness of its use was also influenced by microbial life in the plant root risorfer environment. This is supported by the opinion [9] which states that soil that is sufficiently moist and not dry soil microorganisms grow and function properly, to assist the process of providing and absorbing nutrients. Soil microbial activity in a moist and non-dry soil environment results in a better nitrogen mineralization process.

### 3.4 Support corn straw feed

The results of the calculation of the analysis of the carrying capacity of the feed from corn straw obtained during the study are presented in table 4.

**Table 4:** Calculation results of Corn Straw Feed Carrying Capacity

Parameter	Total Production (ST/Year)
Dry Material	186.42
Crude protein	135.99

Based on table 4 above it can be explained that the dry matter obtained is equal to 186.42 and crude protein obtained by 135.99. Carrying capacity is the ability to provide forage for livestock in the form of natural grass and agricultural waste products from a grazing area. According to [10] carrying capacity is the amount of environmental ability to support animal life, which is expressed by the number of tails per unit of land. The number of animals that can be supported depends on the amount of biomass (plant organic matter) available to the animals. Carrying capacity can be determined by the amount of plant organic matter formed from photosynthesis per unit area and time, which is called primary productivity. The capability of an area can be assessed if the amount of available fodder is greater than the number of livestock accommodated. The carrying capacity index is the result of dividing the potential forage available by the amount of feed needed.

The feed carrying capacity index (IDDP) is used to see the comparison of the availability of forage from food crop waste with the number of ruminant livestock populations in an area. IDDP is obtained from the total production of forage, waste and ram banks available in an area (ST) divided by the number of rumsia livestock population in that area (ST) or the carrying capacity of potential forage divided by the total population [11]. IDDP scores are grouped into three categories, namely low category, medium category, and high category.

Utilization of agricultural waste, namely corn straw, can support the ability of an area to provide forage for livestock. The results of the calculation of the carrying capacity of corn straw in Tanjung Selamat Village, Sunggal District, are 186.5 ST, while the total livestock population is 160 ST. This shows that Sunggal District, Deli Serdang Regency can support the availability of feed from agricultural waste (corn straw).

### 3.5 Analysis of cattle population increase capacity

The results of the calculation of the capacity analysis for increasing the cattle population obtained during the study are presented in table 5 below.

**Table 5:** Calculation Results of Cattle Population Increase Capacity Analysis

Parameter	Total Production (ST* Year)
Real Population	160
Population Increasing Capacity	186.5

Based on table 6 above it can be explained that real population obtained is equal to 160 and K population increase capacity obtained for 186.5. Cattle population increase capacity analysis (KPPTS) is the number of cattle that can be added to an area based on the availability of forages for livestock including forages derived from natural grass and forages derived from food crop waste in an area. According to [12] the use of food crop waste as feed has several obstacles caused by its nutritional value which varies depending on the type of plant, harvest time and post-harvest treatment, besides that plant waste also has high crude fiber so its use is limited. Almost all breeders have not utilized feed processing technologies such as: making silage and making hay when the production of agricultural waste is abundant which can be used during the summer.

The total capacity for increasing the ruminant livestock population is calculated based on the difference between the carrying capacity of forage livestock feed and the number of livestock populations in Tanjung Selamat Village, Sunggal District, Deli Serdang Regency. This is in accordance with the opinion [2] that based on the analysis of the carrying capacity of feed, it can be calculated the increased production capacity of ruminants in an area. Based on the results of the calculation of the capacity to increase the cattle population, it was obtained at 186.5 ST, while data from the cattle population in Tanjung Selamat Village, Sunggal District, Deli Serdang Regency, amounted to 160 ST. This indicates that Tanjung Selamat Village, Sunggal District, Deli Serdang Regency has great potential in increasing and meeting the needs of life and cattle production.

#### 4. Conclusion

Corn straw agricultural waste has not been given optimally to cattle in Tanjung Selamat Village, Sunggal District, Deli Serdang Regency and there is no processing to increase its quality and shelf life. 186.42 ST/harvest/year. Based on the need for crude protein, it can accommodate 135.99 ST/harvest/year. The increase in carrying capacity based on corn straw waste was still able to increase the livestock population to 186.5 ST in Tanjung Selamat Village.

#### Compliance with ethical standards

##### Acknowledgments

The authors would like to thank Animal Husbandry Study Program, Faculty of Sciences and Technology, Pembangunan Panca Budi University.

##### Disclosure of conflict of interest

The authors declare that they have no competing interests.

##### Availability of data and materials

The authors were collected data from this manuscript together.

##### Funding

The research was financed by authors.

#### Authors' contributions

The authors have participated and worked on completing this manuscript and approved the final manuscript

#### 5. References

1. Syamsu JA. Repositioning the Paradigm of Animal Husbandry Development Thoughts, Ideas and Public Enlightenment. Absolut Media, Yogyakarta; c2011.
2. Syamsu JA. Potential Analysis of Food Crop Waste as a Source of Ruminant Animal Feed in South Sulawesi. Dissertation: Graduate School, Bogor Agricultural University, Bogor; c2006.
3. Syamsu JA, Sofyan LA, Mudikdjo K, Gumbira E. Sa'id. Carrying Capacity of Agricultural Waste as a Source of Feed for Ruminants in Indonesia. *Wartazoa*. 2003;13(1):30-37.
4. Sari A. Potential Carrying Capacity of Palawija Plant Waste as Ruminant Animal Feed in Pringsewu District. Thesis. Department of Animal Husbandry, Faculty of Agriculture, University of Lampung, Lampung; c2015.
5. Tangendjaja B, Wina E. Plant Waste and By-Products of the Corn Industry for Feed. Livestock Research Institute. Bogor; c2006.
6. Umela Saiful, Nurfitriyanti B. Carrying Capacity of Corn Straw As Beef Cattle Feed. Department of Agricultural Technology, Gorontalo Polytechnic, Gorontalo; c2016.
7. Sutoro Y, Sulaeman, Iskandar. Cultivation of Corn Plants. Agricultural Research and Development Agency. Food Crops Research and Development Center. Bogor; c1998.
8. Kuzyakov Y. Factor Effecting Rhizosphere Priming Effects. *J Plant dlut Soil Sci*. 2002;165:382-396.
9. Sari A. The Effect of Financial Performance on Changes in Stock Prices by Paying Attention to Company Size. Unpublished Thesis. State University of Surakarta; c2005.
10. Directorate General of Livestock. Technical Guidelines for the Identification and Analysis of the Potential of Livestock Development Areas. Directorate General of Livestock. Agriculture department. Jakarta; c2000.
11. Soetanto. Cassava Processing. Wiyata Jakarta Library and Media Center; c2001.
12. Ance Gunarsih Kartasapoetra. Climatology of Climate Effects on Soil and Plants. *Earth Script: Jakarta*; c2006.