



ISSN: 2456-2912

VET 2023; 8(3): 225-228

© 2023 VET

www.veterinarypaper.com

Received: 19-02-2023

Accepted: 30-03-2023

Minal H Patel

Department of Forest Products
& Utilization, College of
Forestry, Navsari Agricultural
University, Navsari, Gujarat,
India

SS Malek

Department of Silviculture and
Agroforestry, Forest College and
Research Institute, Hyderabad,
Mulugu, Telangana, India

VR Patel

Department of Animal
Nutrition, College of Veterinary
Science & A.H., KU, Navsari,
Gujarat, India

BS Desai

Department of Basic Sciences &
Humanities, College of Forestry,
Navsari Agricultural University,
Navsari, Gujarat, India

SK Jha

Department of Forest Biology
and Tree Improvement, College
of Forestry, Navsari Agricultural
University, Navsari, Gujarat,
India

DP Patel

Department of Natural Resource
Management, College of
Forestry, Navsari Agricultural
University, Navsari, Gujarat,
India

Corresponding Author:

Minal H Patel

Department of Forest Products
& Utilization, College of
Forestry, Navsari Agricultural
University, Navsari, Gujarat,
India

Chemical composition and tannin content in leaves of *Ficus* spp. as top feed for ruminants from the Dang forest of South Gujarat Province, India

Minal H Patel, SS Malek, VR Patel, BS Desai, SK Jha and DP Patel

DOI: <https://doi.org/10.22271/veterinary.2023.v8.i3d.548>

Abstract

An experiment was conducted to estimate proximate composition, fibre fraction and tannin content of leaves amongst 10 spp. of genus *Ficus* L. belonging to family Moraceae from the Dangs Forest in South Gujarat. Results revealed that all parameters varied significantly among different species leaves. Leaves of *F. racemosa* was observed with high ($p < 0.05$) dry matter (DM %) content while *F. arnottiana* leaves was high in organic matter (OM %) amongst. Per cent of crude protein (CP) and ether extract (EE) was significantly high in leaves of *F. benjamina* (11.64%) and *F. virens* (4.23%), respectively. The species *F. asperrima*, *F. arnottiana*, *F. religiosa* and *F. benjamina* were shown to be superior for the parameters like neutral detergent fibre (NDF %), acid detergent fibre (ADF %), acid detergent lignin (ADL %), and cell content. Tannin content was low in *F. benghalensis* and *F. benjamina* while highest in *F. arnottiana* followed by *F. amplissima*. Among different studied species *F. benghalensis* and *F. benjamina* found most suitable top feed for ruminant feeding.

Keywords: Proximate parameters, fibre fraction, tannins, ficus, moraceae, ruminant

Introduction

For livestock animals, fodder serves as their main and basic source of nutrition and is essential to their growth, development, and general well-being. Species, age, gender, and production goals all affect the nutritional needs of livestock animals^[10]. To meet the nutritional requirement of increasing livestock population, it is necessary both to increase the productivity and to use available resources more efficiently. The leaves of tree considered nutritious feed due to their high proteins, vitamins and minerals^[17]. Tree leaves play an important role in the nutrition of grazing animals in area where few or no alternatives are available^[20].

For instance, in order to support milk production, dairy cows need a diet high in calories and protein, whereas beef cattle need a diet high in fibre to support healthy rumen function^[9]. Because of the secondary plant chemicals (tannins) found in tree leaves, which allow ruminants to acquire larger quantities of dietary protein at post rumen for digestion and absorption, trees are employed as forages as sources of protein and energy for small ruminants^[5]. For livestock animals to obtain enough nutrients from their food, it is essential to understand and determine the nutritional makeup of various types of fodder. For this, it is crucial to comprehend the basic nutrient composition of fodder, including its fiber fraction and tannin content.

The Moraceae family includes a wide range of trees and shrubs collectively known as *Ficus* spp., or simply figs. These plants are widespread in tropical and subtropical areas and are valued for their decorative qualities, edible fruits, and ecological significance^[3,4]. But one of the less well-known advantages of *Ficus* species is their potential as a source of cattle fodder. *Ficus* spp. leaves and fruits are extremely nutrient-dense, loaded with protein and minerals, and have long been utilised as animal feed in many cultures^[28]. Exploring the potential of *Ficus* spp. as a sustainable and cost-effective fodder source can help to promote livestock farming and conservation initiatives in a variety of areas. We can increase animal health, productivity, and welfare while lowering the environmental impact of animal husbandry by optimizing the nutritional content of livestock feeds.

Materials and Methodology

Tree leaves have traditionally been used as sources of fodder since they are a natural component of ruminant diets. For tiny ruminants, tree fodders constitute a significant source of nutrition. Tree fodders can supplement low-quality grasses since they provide more protein and minerals than grasses [5]. To ascertain the nutritional content of leaves from locally accessible fodder tree species, a field survey experiment was carried out in Dang district of South Gujarat. The Dang comes under the AES-III, South Gujarat Heavy Rainfall Zone with a hilly terrain, is governed by dryness except during the rainy season. It consists of mixed tree growth among which teak is the predominant species in most parts of the area. The forest is continuous in all parts except at the places where the slopes are too steep to retain the soil and where biotic interference has cleared or opened it out.

Leaf fodder of ten different *Ficus* species viz., (*Ficus asperrima* Roxb., *F. racemosa* L., *F. virens* Ait., *F. benghalensis* L., *F. religiosa* L., *F. hispida* Vahl., *F. amplissima* Sm., *F. rumphii*, *F. benjamina* L., *F. arnottiana* Miq.) was collected from Dang Forest, Gujarat in the Summer season (April-May) of year 2022-23. Proximate parameters such as moisture content (MC), dry matter (DM), organic matter (OM), crude protein (CP), crude fiber (CF), ether extract (EE), total ash (TA) and nitrogen free extract (NFE) along with fibre fraction such as neutral detergent fibre (NDF), acid detergent fibre (ADF), acid detergent lignin (ADL), cell content, cellulose and hemi-cellulose content of collected leaves were analyzed as per Van soest *et al.*, (1991) [33] and AOAC (2016) [1] in three replication and the further data were analysed using CRD analysis [26, 14]. The tannin content of leaves was estimated though by using Folin

Ciocalteu assay method [21].

Result and Discussion

The proximate composition and fiber fractions of different *Ficus* tree leaves were revealed significant variations ($P \leq 0.05$) among different species (Table 1). The maximum, OM% (92.95) and CF% (27.91) were significantly higher in leaves of *F. arnottiana* however, DM % was found higher in *F. racemosa* (47.27%). CP (%) content was high ($p < 0.05$) in *F. benjamina* (11.64%), which was comparable with *F. benghalensis* (11.41%) and *F. racemosa* (11.29%). EE (%) was found in the range of 1.23 to 4.23, which is better to that of conventional fodder like maize (1.79%) and sorghum (1.94%) [18, 29].

Generally, CP content typically provides a reliable indicator of their nutritional status, however it varies widely and similar research was conducted by Osagie and Aguebor-Ogie (2020) [24] on the three *Ficus* species (*F. thonningii*, *F. carica*, and *F. exasperata*) and reported the CP range (7.63-15.76%), which validates and illustrates the range (8.51-11.64%) of our results.

Similarly, in our study we reported significant variations ($p > 0.05$) in proximate composition like OM, CF, CP, EE and the ash, NFE and crude fibre concentration among the ten *Ficus* species studied in our study which validates our finding and there are also comparable to values previously reported for Osowe *et al.*, 2021 [25] in different species of *Ficus*; *F. racemosa* [11, 30, 34, 7] and *F. benghalensis* [19, 30, 27, 12]. Chitra and Balasubramanian 2016 [6] also reported the different proximate composition and fiber fraction in the five *Albizia* species which is also supports our results.

Table 1: Proximate content in the different *Ficus* spp. from the Dang Forest of South Gujarat Province, India

#	MC (%)	DM (%)	OM (%)	N (%)	CP (%)	EE (%)	CF (%)	AC (%)	NFE (%)
1	61.15 ^d	38.85 ^d	81.73 ^e	1.62 ^c	10.12 ^c	1.35 ^f	26.39 ^a	18.27 ^a	43.86 ^d
2	71.19 ^a	28.81 ^g	92.95 ^a	1.37 ^e	8.58 ^e	1.74 ^e	27.91 ^a	7.05 ^e	54.72 ^{ab}
3	67.70 ^b	32.30 ^f	87.37 ^b	1.77 ^b	11.07 ^b	2.34 ^d	17.99 ^{cd}	12.63 ^d	55.97 ^{ab}
4	59.15 ^e	40.85 ^c	85.14 ^{bc}	1.83 ^{ab}	11.41 ^{ab}	2.23 ^d	20.78 ^b	14.86 ^{cd}	50.72 ^c
5	65.03 ^c	34.97 ^e	83.25 ^{cde}	1.86 ^a	11.64 ^a	1.23 ^f	16.72 ^{cde}	16.75 ^{abc}	53.67 ^{bc}
6	67.71 ^b	32.28 ^f	84.00 ^{cd}	1.62 ^c	10.10 ^c	1.36 ^f	15.15 ^c	16.00 ^{bc}	57.40 ^a
7	52.73 ^g	47.27 ^a	86.66 ^b	1.80 ^{ab}	11.29 ^{ab}	3.86 ^b	15.79 ^{de}	13.34 ^d	55.72 ^{ab}
8	63.78 ^c	36.22 ^e	91.77 ^a	1.52 ^d	9.51 ^d	3.58 ^c	25.68 ^a	8.23 ^e	53.00 ^{bc}
9	57.53 ^f	42.47 ^b	82.28 ^{de}	1.36 ^e	8.51 ^e	3.71 ^{bc}	12.10 ^f	17.72 ^{ab}	57.95 ^a
10	56.52 ^f	43.48 ^b	85.42 ^{bc}	1.40 ^e	8.78 ^e	4.23 ^a	19.17 ^{bc}	14.58 ^{cd}	53.25 ^{bc}
S.Em	0.53	0.53	0.77	0.03	0.18	0.07	0.92	0.77	1.21
CV%	1.46	2.41	1.55	3.20	3.14	4.19	8.06	9.56	3.91

[MC: Moisture content (%), OM: Organic Matter (%), N: Nitrogen (%), CP: Crude Protein (%), EE: Ether extract (%), CF: Crude fibre (%), AC: Ash Content (%), NFE: (Nitrogen free extract. %), 1: *F. amplissima*, 2: *F. arnottiana*, 3: *F. asperrima*, 4: *F. benghalensis*, 5: *F. benjamina*, 6: *F. hispida*, 7: *F. racemosa*, 8: *F. religiosa*, 9: *F. rumphii*, 10: *F. virens*. {means with same and different superscript letter in the same column indicate significant and insignificant difference ($p < 0.05$).}]

In ruminant system, nutrient digestibility is affected by content of fiber and its fractions which is shown in Table 2. Highest CF (%) was observed in *F. arnottiana* (27.91%), which is also found high in ADF (52.40%). Leaves of *F. asperrima* was found significantly high for NDF (72.42%) and ADL (36.43%) content. Cellulose and Hemi cellulose being considered as major polysaccharides available from fibrous portion of roughages. In the present study, highest cellulose and hemicellulose was found in *F. arnottiana* (17.64%) and *F. asperrima* (21.55%), respectively. The ADL portion contributed by lignin and total ash, which are

indigestible and reduce the digestibility of fodder. Here, highest ($p < 0.05$) ADL was observed in *F. religiosa* (38.09%) followed by *F. asperrima* (36.43%).

Results of fiber fraction are supported by the Niranjana, 2007 [23], Dhungana *et al.*, 2012 [7], Nayak and Basak, 2015 [22] for the *F. benghalensis*. However, values for NDF (42.43-72.72%) was found higher than Ramachandran *et al.*, 2015 [27] and Gaikwad *et al.*, 2017 [12] for *F. religiosa*. Similarly, proximate parameter of *F. virens* reported in the study was in corroboration with of Khan 2011 [16] and Niranjana 2007 [23].

Table 2: Fiber fraction and Tannins content in the different *Ficus* spp. from the Dang Forest of South Gujarat Province, India

#	NDF (%)	ADF (%)	ADL (%)	Cellulose (%)	Hemi-cellulose (%)	CC (%)	Tannins (%)
1	46.94 ^f	29.45 ^e	12.79 ^f	16.66 ^a	17.49 ^c	53.06 ^b	27.37 ^b
2	64.59 ^b	52.40 ^a	34.76 ^b	17.64 ^a	12.20 ^f	35.41 ^f	38.56 ^a
3	72.42 ^a	50.87 ^a	36.43 ^a	14.43 ^b	21.55 ^a	27.58 ^g	11.81 ^{cd}
4	52.96 ^d	40.29 ^b	34.06 ^{bc}	6.23 ^c	12.66 ^{ef}	47.04 ^d	8.10 ^g
5	42.43 ^g	29.34 ^e	24.53 ^e	4.80 ^{cd}	13.09 ^{ef}	57.57 ^a	9.46 ^{fg}
6	46.83 ^f	33.39 ^d	30.01 ^d	3.38 ^{de}	13.44 ^e	53.17 ^b	11.00 ^{de}
7	49.24 ^e	36.67 ^c	32.46 ^c	4.21 ^{de}	12.58 ^{ef}	50.76 ^c	11.85 ^{cd}
8	55.49 ^c	40.84 ^b	38.09 ^a	2.76 ^e	14.65 ^d	44.51 ^e	12.61 ^c
9	51.46 ^d	32.42 ^d	29.56 ^d	2.86 ^e	19.05 ^b	48.54 ^d	10.19 ^{ef}
10	57.33 ^c	40.18 ^b	33.96 ^{bc}	6.22 ^c	17.15 ^c	42.67 ^e	12.31 ^{cd}
S.Em	0.73	0.80	0.57	0.65	0.31	0.73	0.47
CV%	2.34	3.60	3.19	14.22	3.53	2.75	5.27

[NDF: (Neutral detergent fiber %), ADF: (Acid detergent fiber %), ADL: (Acid Detergent Lignin %), C: Cellulose (%), HC: Hemi-Cellulose (%), CC: Cell content (%), DM: Dry Matter (%), 1: *F. amplissima*, 2: *F. arnottiana*, 3: *F. asperrima*, 4: *F. benghalensis*, 5: *F. benjamina*, 6: *F. hispida*, 7: *F. racemosa*, 8: *F. religiosa*, 9: *F. rumphii*, 10: *F. virens*. {means with same and different superscript letter in the same column indicate significant and insignificant difference ($p < 0.05$).}]

The common *Ficus* trees fodder is quite usual among the cattle and small ruminants. However, no information is available on the nutritive value for few of the selected species. It might be the first attempt to analysed *Ficus asperrima*, *F. amplissima*, *F. rumphii*, *F. benjamina* and *F. arnottiana* for their proximate and fiber fraction composition.

The tannin content of fodder plants, which has a substantial impact on digestibility and nutrient availability, is one element that affects the nutritional value of the plants [15]. The majority of tree leaves include different types of tannins as an antinutritional substance that necessitates appropriate measures while choosing the species of tree leaves [5].




Fig 1: Tannin (%) content in the different *Ficus* spp. from the Dang Forest of South Gujarat Province, India. (1: *F. amplissima*, 2: *F. arnottiana*, 3: *F. asperrima*, 4: *F. benghalensis*, 5: *F. benjamina*, 6: *F. hispida*, 7: *F. racemosa*, 8: *F. religiosa*, 9: *F. rumphii*, 10: *F. virens*.)

In present investigation of *Ficus* spp., the tannin was found in the range of 8.10-38.56%. Maximum mean tannin content was observed in the FAR (*Ficus arnottiana*) leaves 38.56% followed by FAM (*F. amplissima*) leaves 27.37%. However, the minimum mean tannin content was observed in FB (*Ficus benghalensis*) 8.10% and shown in the Table 2 and Fig. 1. Similar findings were reported by Fernandes *et al.*, (2007) [11] and Sumi *et al.*, (2021) [32], they reported that the tannin levels in *Ficus racemosa* was 2.9% and 19.72 mg GAE/g of dry extract, respectively by Folin Ciocalteu method. Numerous reports that have shown the reduction of enteric methane from ruminants due to inclusion of tannin rich browses because the tannins have anti-methanogenic activity [13] and ranged the total tannin concentration between 32.4 to 209.1 g/kg DM. Similarly, Saxena *et al.*, 2013 [31] reported 99.55 mg/gm, 9.95mg/gm, 54.96 mg/gm, and 57.4869 mg/gm of tannin for Harde, Arjuna, Baheda and Ashoka respectively.

Conclusion

The present study on nutritional evaluation of leaves from different *Ficus* species have been revealed that leaves are good source of nutrients with tolerable amount of ADL as well as tannin content and can be used as a supplemental feed in livestock.

Acknowledgement

Acknowledging the Department of Animal Nutrition, Vanbandhu College of Veterinary Science & Animal Husbandry, Navsari Campus, KAU, Gandhinagar for providing the necessary facilities for the analysis.

References

1. AOAC. Official methods of Analysis, (19th Ed). Association of Official Analytical Chemist. Washington DC; c2016.
2. Khan WA. *Ficus palmata* a nutraceutical under took advantage of the product. Int. J Agric. Nutr. 2021;3(2):14-18. DOI: 10.33545/26646064.2021.v3.i2a.67
3. Berg CC, Corner EJ, Noteboom HP. Moraceae (*Ficus*). Flora Malesiana, Series I (Seed plants). Leiden: National Herbarium of the Netherlands; 2005;17(2):01-70.
4. Chaudhary LB, Sudhakar JV, Kumar A, Bajpai O, Tiwari R, Murthy GV. Synopsis of the genus *Ficus* L. (Moraceae) in India. Taiwan. 2012;57(2):193-216.
5. Cheema UB, Younas M, Sultan JL, Virk MR, Tariq M, Waheed A, *et al.* Fodder tree leaves: an alternative source

- of livestock feeding. *Advances in Agricultural Biotechnology*. 2011;2:22-33.
6. Chitra P, Balasubramanian A. A study on chemical composition and nutritive value of *Albizia* tree leaves as a livestock feed. *International Journal of Science, Environment and Technology*. 2016;5(6):4638-4642.
 7. Damame SV, Gore SB, Sonone AH. Nutritional Composition of Common Fodder Tree Leaves from Ahmednagar District of Maharashtra State. *BIOINFOLET-A Quarterly Journal of Life Sciences*. 2016;13(2b):394-397.
 8. Dhungana S, Tripathee HP, Puri L, Timilsina YP, Devkota KP. Nutritional analysis of locally preferred fodder trees of middle hills of Nepal: A Case Study from Hemja VDC, Kaski District. *Nepal Journal of Science and Technology*. 2012;13(2):39-44.
 9. Erickson PS, Kalscheur KF. Nutrition and feeding of dairy cattle. *Animal Agriculture*; c2020. p. 157-180.
 10. FAO. *World Livestock: Transforming the livestock sector through the Sustainable Development Goals*. Rome; c2018. p. 222. <https://doi.org/10.4060/cal201en>. Licence: CC BY-NC-SA 3.0 IGO.
 11. Fernandes AP, Shintre PM, Fulpagare YG. Nutritive value of top feeds for goats. *Indian Journal of Animal Nutrition*. 2007;24(1):40-43.
 12. Gaikwad US, Pawar AB, Kadlag AD. Nutritional status of fodder tree leaves and shrubs of scarcity zone of Maharashtra. *Advances in Life Sciences*. 2017;7(1):11-14.
 13. Gameda BS, Hassen A. Effect of tannin and species variation on *in vitro* digestibility, gas, and methane production of tropical browse plants. *Asian-Australasian Journal of Animal Sciences*. 2015;28(2):188-199.
 14. Gopinath PP, Parsad R, Joseph B, Adarsh VS. Grapes Agri1: collection of shiny apps for data analysis in agriculture. *Journal of Open-Source Software*. 2021;6(63):3437. <https://doi.org/10.21105/joss.03437>.
 15. Hassan ZM, Manyelo TG, Selaledi L, Mabelebele M. The effects of tannins in monogastric animals with special reference to alternative feed ingredients. *Molecules*. 2020;25(20):4680. Doi: 10.3390/molecules25204680.
 16. Khan N, Barman K, Rastogi A, Sharma RK, Yattoo MA. Chemical composition, tannin fractionation and protein binding affinity of some top foliages. *Indian Journal of Animal Nutrition*. 2011;28(4):421-426.
 17. Lata M, Mondal BC. Uses of tree leaves as alternative feed resources for ruminant animals. *Journal of Entomology and Zoology Studies*. 2021;9(1):2239-2247.
 18. Telgote MV, Chavan SD, Shelke RR, Nage SP, Bidwe KU. Comparative study on chemical composition of different feeding ingredients with hydroponically grown and conventionally grown green maize fodder. *Biological Forum – An International Journal*. 2022;14(3):656-659.
 19. Maw NN, San Mu K, Aung A, Htun MT. Preliminary report on nutritive value of some tree foliages. In *Conference on International Agricultural Research for Development*; c2011. p. 11-13.
 20. Meuret M, Boza J, Narjisse H, Nastis A. Evaluation and utilization of rangeland feeds by goats. *Goat Nutrition*. 1991;46:160.
 21. Mohammed S, Manan FA. Analysis of total phenolics, tannins and flavonoids from *Moringa oleifera* seed extract. *Journal of Chemical and Pharmaceutical Research*. 2015;7(1):132-135.
 22. Nayak J, Basak UC. Analysis of some nutritional properties in eight wild edible fruits of Odisha, India. *International Journal of Current Science Research and Review*. 2015;14(1):55-62.
 23. Niranjana PS, Srivastava V, Verma DN. Nutritional evaluation of peepal leaves in barbari kids. *Indian Journal of Animal Nutrition*. 2007;24(2):128-129.
 24. Osagie OA, Aguebord-ogie BN. Proximate Analysis, Phytochemical Screening and Evaluation of Acute and Sub-chronic Toxicity of Methanol Extract of *Ficus exasperata*. *Federation of American Societies for Experimental Biology Journal*. 2020;34(S1):1-1. <https://doi.org/10.1096/fasebj.2020.34.s1.00304>
 25. Osowe CO, Olowu OP, Adu OA, Oloruntola OD, Chineke CA. Proximate and mineral composition, phytochemical analysis, and antioxidant activity of fig trees (*Ficus* spp.) leaf powder. *Asian Journal of Biochemistry, Genetics and Molecular Biology*. 2021;9(1):19-29.
 26. Panse VG, Sukhatme PV. *Statistical methods for agricultural workers*. ICAR, New Delhi; c1967.
 27. Ramachandran M, Bharathidhasan A, Balakrishnan V. Nutrient composition, *in vitro* true digestibility (IVTD) and methane production potential of fodder tree leaves. *Indian Journal of Animal Sciences*. 2015;85(5):494-497.
 28. Rasool IFu, Aziz A, Khalid W, Koraqi H, Siddiqui SA, AL-Farga A, *et al*. Industrial Application and Health Prospective of Fig (*Ficus carica*) By-Products. *Molecules*. 2023;28(3):960. <https://doi.org/10.3390/molecules28030960>.
 29. Roy A, Roy PS, Mandal N, Kumar S. Evaluation of Chemical Composition of Fifteen Commonly used Ruminants Feed Ingredients locally available in Murshidabad District, West Bengal: Evaluation of chemical composition of fifteen commonly used ruminants feed ingredients locally available in Murshidabad District, West Bengal. *Indian Journal of Agricultural Research*. 2019;6(1):135-138.
 30. Sahoo A, Singh B, Bhat TK. Effect of tannins on *in vitro* ruminal protein degradability of various tree forages. *Livestock Research for Rural Development*. 2010;22(7):119.
 31. Saxena VA, Mishra G, Saxena A, Vishwakarma KR. A comparative study on quantitative estimation of tannins in *Terminalia chebula*, *Terminalia belerica*, *Terminalia arjuna* and *Saraca indica* using spectrophotometer. *Asian Journal of Pharmaceutical and Clinical Research*. 2013;6(3):148-149.
 32. Sumi SA, Siraj MA, Hossain A, Mia MS, Afrin S, Rahman MM, *et al*. Pharmacological activities of *Ficus racemosa* and analysis of its major bioactive polyphenols by HPLC-DAD. *Highlights on Medicine and Medical Science*. 2021;5:68-84.
 33. Van Soest PV, Robertson JB, Lewis BA. Methods for dietary fiber, neutral detergent fiber, and non-starch polysaccharides in relation to animal nutrition. *Journal of Dairy Science*. 1991;74(10):3583-97.
 34. Bakshi MP, Wadhwa M. Comparative nutritional evaluation of forest tree leaves and wild grasses of semi hilly arid zone in Punjab. *Indian Journal of Animal Sciences*. 2012;82(8):918.